arXiv:1005.4736v4 [math.GR] 20 Jul 2010

Multielement order separability in free products of groups [1]

Vladimir V. Yedynak

Abstract

We introduce the notion of multielement order separability and study this property for free groups and free products.

Keywords: free products, residual properties.
MSC: 20E26, 20E06.

1 Introduction.

Definition 1. We say that a group G is *n-element order separable* if, for each elements $u_1, ..., u_n$ of G such that u_i and $u_j^{\pm 1}$ are not conjugate for $i \neq j$, there exists a homomorphism of G onto a finite group such that the orders of the images of $u_1, ..., u_n$ are pairwise different.

It is easy to see that, if a free product of two groups is n-element order separable, then the free factors are k-element order separable for each $k \leqslant n$.

The converse is not true for $n > 2$. For example, the infinite dihedral group $\langle a \rangle_2 * \langle b \rangle_2$ is not 3-element order separable.

So, we require additionally each two nontrivial elements from different free factors not to have equal finite orders. It is known that free groups are 2-element order separable (Equations over groups, quasivarieties, and a residual property of a free group, *Journal of group theory* **2**: 319-327), and, moreover, this property is inherited by free products (Separability with respect to order, *Vestnik Mosk. Univ. Ser. I Mat. Mekh.* **3**: 56-58).

It was proven that free metabelian groups are 2-element order separable, but this property fails for free 3-step solvable groups (Elements with the same normal closure in a metabelian group, *The Quarterly Journal of Mathematics* **58(1)**: 23-29).

Theorem 1. *Any absolutely free group is n-element order separable for each n.*

Theorem 2. *Suppose that groups A and B are periodic and n-element order separable for each n. If $|a| \neq |b|$ for any $a \in A \setminus \{e\}, b \in B \setminus \{e\}$, then the group $A * B$ is also n-element order separable for each n.*

Theorem 3. *If groups A and B are n-element order separable for each n and $|a| \neq |b|$ for any $a \in A \setminus \{e\}, b \in B \setminus \{e\}$, such that $|a| < \infty, |b| < \infty$, then the free product $G = A * B$ is 3-element order separable.*

---

[1] This work was supported by the Russian Foundation for Basic Research, project no. 05-01-00895

2 Notations and definitions.

Suppose that $G = A * B$, where A and B are finite groups. Consider an oriented graph Γ with the following properties:

(1) For each vertex p and each $c \in A \cup B \setminus \{1\}$, there are exactly one oriented edge labelled by c and starting at p, exactly one oriented edge labelled by c and ending at p.

(2) For each vertex p, the maximal oriented connected subgraph $A(p)$ containing p, whose edges are labelled by elements of the group A, is the Cayley graph of the group A with respect to the set of generators A. Similarly, for each vertex p, the maximal oriented connected subgraph $B(p)$ containing p, whose edges are labelled by elements from the group B, is the Cayley graph of the group B with respect to the set of generators B.

If e is an edge of the graph Γ, then $\mathrm{Lab}(e)$ denotes the label of e. If $s = e_1...e_n$ is a path in Γ, then $\mathrm{Lab}(s) = \mathrm{Lab}(e_1)...\mathrm{Lab}(e_n)$ denotes the label of the path s, $\alpha(s) = \alpha(e_1)$ is the beginning of e_1, $\omega(s) = \omega(e_n)$ is the end of e_n.

Definition 2. *Let $x \in G$ be a cyclically reduced element not belonging to $A \cup B$. We say that a closed path $e_1...e_n$ whose label equals x^k is an x-cycle if Lab($e_{il+1}...e_{il+l}$) is the normal form of the element x, where l is the length of the word x, i is an arbitrary positive integer, and indices are modulo n. The number k is called the length of this x-cycle.*

Definition 3. *If two different edges of a cycle lies in the same subgraph $A(p)$ or $B(p)$, we say that this cycle has close edges.*

Let Γ be a graph with Properties (1), (2). Take a set of vertices S in the graph Γ. Consider an arbitrary function $f : S \to \{A, B\}$. Consider t copies $\Delta_1, ..., \Delta_t$ of Γ. Let $v_i \in \Delta_i$ be the vertex corresponding to a vertex $v \in \Gamma$, let $q_i \in \Delta_i$ be the edge corresponding to an edge $q \in \Gamma$ and let S_i be the set of vertices of Δ_i corresponding to the set S. For each $i \in \{1, \dots, t\}$, for each edge q starting or ending at a vertex from S and each $p \in S$, we delete edges labelled by the elements of the group $f(p)$ whose end points coincide with p_i, if q_i starts at p_i, ends at v_i and $\mathrm{Lab}(q_i) \in f(p)$ connect the vertices p_i and v_{i+1} (subscripts modulo t) by an oriented edge starting at p_i whose labell equals $\mathrm{Lab}(q_i)$, if q_i starts at v_i, ends at p_i and $\mathrm{Lab}(q_i) \in f(p)$ connect the vertices p_i and v_{i+1} (subscripts modulo t) by an oriented edge starting at v_{i+1} whose labell equals $\mathrm{Lab}(q_i)$. Let $S = \{s_1, \dots, s_k\}$. The obtained graph $\gamma_t(\Gamma; s_1, \dots, s_k; f(s_1), \dots, f(s_k))$ satisfies Properties (1), (2).

The group G acts on the right on the set of vertices of the graph Γ with Properties (1), (2) by the following way. If $a \in A$ and p is a vertex of Γ then there exists exactly one edge h starting at p and labelled by a. Put $p \cdot a = \omega(h)$. (Here and in what follows, $\alpha(h)$ and $\omega(h)$ denote the start and end points of an oriented edge h.) The action of group B is defined similarly.

In what follows, having a homomorphism φ of the group $A * B$ we always assume that the Cayley graph of the group $\varphi(A * B)$ is constructed with respect to the set of generators $\varphi(A \cup B)$.

3 Auxiliary Lemmas.

Lemma 1. *If $G = A * B$, where the groups A and B are finite, $w_1, ..., w_k \in C \backslash e$ (throughout this paper, C denotes the Cartesian subgroup of the free product of groups A and B), and p is a prime number, N is a positive integer, then there exists a homomorphism φ of G onto a finite group such that, for each i, $\varphi(w_i)$ is a p-element and $|\varphi(w_i)| > p^N$.*

Proof.

Since C is a free group, there exists a homomorphism φ_1 of C onto a finite p-group such that $\varphi_1(w_i^j) \neq 1$ for $i = 1, ..., k, j = 1, ..., N$ (Foundations of group theory, *Nauka*, theorem 14.2.2). Put $N = \text{Ker } \varphi_1$. Consider the subgroup $N' = \cap_{g \in G}(g^{-1}Ng)$. Then $N' \lhd G, |G : N'| < \infty$ and the order of the image of w_i in the group G/N' is a power of p which is greater than N for each i. The natural homomorphism $\varphi : G \longrightarrow G/N'$ is as required.

Lemma 2. *Suppose that $G = A * B$, where A and B are finite groups. Let S be the finite set of elements, such that for any $u \in S$ the following conditions are satisfied: u is a nonunit cyclically reduced element of the Cartesian subgroup C, furthermore $u = u'^m$, where u' is not a proper power and if $m > 1$ then for each $m' \neq m$ such that $m'|m$ $u'^{m'} \notin C$. Then for any prime p there exists a homomorphism φ of G onto a finite group such that for each $u \in S$ $\varphi(u)$ is a nonunit p-element, and all u-cycles in the Cayley graph of $\varphi(G)$ have no close edges.*

Proof.

Consider a prime p. Fix an element u of S. Put $u' = y_1...y_n$ — the normal form of u'. Fix μ, ν such that $1 \leqslant \mu, \nu \leqslant n$. Fix also $z \in A \cup B$. We consider that if $z = 1$ then $\mu \neq \nu + 1$ and $\nu - \mu < n - 1$.

If $\mu > 1$ then $\mu' = y_\mu...y_n$ otherwise $\mu' = 1$. If $\nu < n$ then $\nu' = y_1...y_\nu$ otherwise $\nu' = 1$. Consider an element $\chi_{z,\mu,\nu} = \nu' z \mu'$. It is easy to see that $\chi_{z,\mu,\nu} \notin \langle u' \rangle$. Let us to prove that there exists a homomorphism $\varphi_{z,\mu,\nu}$ of G onto a finite group such that for each $w \in S$ $\varphi_{z,\mu,\nu}(w)$ — nonunit p-element, $\varphi_{z,\mu,\nu}(\chi_{z,\mu,\nu}) \notin \langle \varphi_{z,\mu,\nu}(u') \rangle$, and $|\varphi_{z,\mu,\nu}(u')| = m|\varphi_{z,\mu,\nu}(u)|$. Lemma 1 shows that there exists a homomorphism φ_1 of G onto a finite group such that $\varphi_1(w)$ is a nonunit p-element for each $w \in S$. It is obvious that $|\varphi_1(u')| = m'|\varphi_1(u)|$, where m' is a divisor of m. Since $u'^m \in C$ and $u'^{m'|\varphi_1(u)|} \in \text{Ker } \varphi_1 \subset C$ then $u'^{m'} \in C$. Hence we deduce that $m' = m$ and $|\varphi_1(u')| = m|\varphi_1(u)|$. We may assume that $\varphi_1(\chi_{z,\mu,\nu}) \in \langle \varphi_1(u') \rangle$.

Let Γ be the Cayley graph of the group $\varphi_1(G)$. Consider an arbitrary vertex r in the graph Γ and a path $L = f_1...f_t$ starting at r such that $\text{Lab}(L)$ is the normal form of $\chi_{z,\mu,\nu}$. Suppose that $\omega(L) = q$. By the assumption, there exists a u'-cycle $S = e_1...e_k$ starting at q, and there exists an edge e_l such that $\omega(e_l) = \alpha(L)$ and $\text{Lab}(e_1...e_l) = u'^v$. Let $\Delta_1, ..., \Delta_p$ be copies of the graph Γ and let e_i^j be the edge of Δ_j corresponding to the edge e_i in Γ. Let $\chi_{z,\mu,\nu} = x_1...x_s$, be the normal form of $\chi_{z,\mu,\nu}$.

If $y_n \in A$ then D denotes A and F denotes B, if $y_n \in B$ then D denotes B, and F denotes A.

If x_1 and y_n belong to different free factors, we construct the graph $\Delta = \gamma_p(\Gamma; r, \omega(e_{l+1}); D, D)$ from the copies Δ_i.

If x_1 and y_n belong to the same free factor, then we construct the graph $\Delta = \gamma_p(\Gamma; r, \alpha(e_l); F, F)$ from the copies Δ_i.

It is obvious that the length of each u'-cycle equals either $|\varphi_1(u')|$ or $p|\varphi_1(u')|$ in Δ. Besides, the length of each w-cycle equals either $|\varphi_1(w)|$ or $p|\varphi_1(w)|$ in Δ for any $w \in S$. The group G acts on the right on the set of vertices of Δ. Besides, if r_i is a vertex of Δ_i corresponding to the vertex r of Γ, then $r_i \cdot \chi_{z,\mu,\nu} u^l \neq r_i$ in Δ for any integer l. Thus, we found the required homomorphism $\varphi_{z,\mu,\nu}$. Besides, $|\varphi_{z,\mu,\nu}(u')| = m|\varphi_{z,\mu,\nu}(u)|$ and $\varphi_{z,\mu,\nu}(w)$ — nonunit p-element for any $w \in S$.

Consider a homomorphism $\varphi_u : G \to \prod_{z,\mu,\nu}(\varphi_{z,\mu,\nu}(G))$ defined by the formula: $g \mapsto \prod_{z,\mu,\nu} \varphi_{z,\mu,\nu}(g)$, where $z \in A \cup B, 1 \leqslant \mu, \nu \leqslant n$ and if $z = 1$ then $\mu \neq \nu + 1, \nu - \mu < n - 1$. In the Cayley graph $Cay(\varphi_u(G))$ of the group $\varphi_u(G)$ all u'-cycles have no close edges and, $\varphi_u(w)$ — is a nonunit p-element for any $w \in S$. Furthermore $|\varphi_u(u')|$ is the less common multiple of numbers $m|\varphi_{z,\mu,\nu}(u)|$ that is $m|\varphi_u(u)|$. Hence all u-cycles in $Cay(\varphi_u(G))$ also have no close edges. It is obvious that the homomorphism $\varphi : G \to \prod_{u \in S}(\varphi_u(G))$ defined by the formula $g \mapsto \prod_{u \in S} \varphi_u(g)$ is as required.

Lemma 3. *Suppose that $G = A * B$, where A and B are finite groups, $u_1, ..., u_m$ are cyclically reduced elements of the Cartesian subgroup such that each two of them do not belong to conjugate cyclic subgroups, π is a finite set of prime numbers. Besides $u_i = u_i'^{m_i}$, where u_i' is not a proper power and if $m_i > 1$ then for each $m_i' \neq m_i$ such that $m_i'|m_i$ $u_i'^{m_i'} \notin C$. Then there exists a homomorphism φ of G onto a finite group such that the images of $u_1, ..., u_m$ have pairwise different orders and $p \nmid |\varphi(u_i)|$ for each i and for each $p \in \pi$.*

Proof.

Let us prove by the induction on m. If $m = 1$, the assertion follows from Lemma 1.

Let us prove the following statement. For elements $u_1, ..., u_m$, there exists a homomorphism φ_1 of G onto a finite group such that the images of $u_1, ..., u_m$ are nonidentity p-elements, $p \notin \pi$; in addition $\{u_1, ..., u_m\} = \alpha \cup \beta$, where α and β are nonempty sets with empty intersection, moreover, for each $u_i \in \alpha$ and $u_j \in \beta$, we have $|\varphi_1(u_i)| > |\varphi_1(u_j)|$.

According to Lemma 2, there exists a homomorphism ψ of G onto a finite group such that $\psi(u_1), ..., \psi(u_m)$ are nonidentity p-elements, where p is a prime and $p \notin \pi$. Furthermore, if Γ is the Cayley graph of the group $\psi(G)$, then, in the graph Γ, all u_i-cycles have no close edges for each i. We may assume that $|\psi(u_1)| = ... = |\psi(u_m)|$. Consider some set of nonnegative integers k. For each k from this set, we construct a graph Γ_k satisfying (1), (2), and the following properties:

(3) the lengths of maximal u_i-cycles coincide for $i = 1, \ldots, m$;

(4) for each i, the length of each u_i-cycle divides the length of a maximal u_i-cycle;

(5) u_i-cycles have no close edges;

(6) if $k > 0$, then there exists a path R_k of length k lying in some maximal u_1-cycle and in all maximal u_j-cycles for each $j = 2, ..., m$;

Properties (1), (2) and (3)–(6) are obviously true for the graph $\Gamma_0 = \Gamma$.

Having a graph Γ_k, we construct a new graph Γ^*_{k+1} satisfying Properties (1), (2) and (4), (5). Hence we shall deduce that this graph satisfies Properties (3), (6) so we put $\Gamma_{k+1} = \Gamma^*_{k+1}$; otherwise, owing to the Property (5), we will prove that the least common multiple of the lengths of all u_1-cycles and the least common multiple of the lengths of all u_j-cycles in Γ^*_{k+1} do not coincide for some j.

Let n be the length of a maximal u_1-cycle in Γ_k. Consider a u_1-cycle S in Γ_k of length n. Suppose that if $k > 0$ then S contains R_k.

Suppose that $k = 0$. Let $S = f_1 ... f_k$. Put $s = \omega(f_1)$ and $t = \omega(f_2)$. If $\mathrm{Lab}(f_1) \in A$ then D denotes A and F denotes B, if $\mathrm{Lab}(f_1) \in B$ then D denotes B and F denotes A. Consider the graph $\Gamma^*_1 = \gamma_{n^2}(\gamma_n(\Gamma_0; s; D); t_{2,1}; F)$. Let $t_2 \in \gamma_n(\Gamma_0; s; D)$ be the vertex of the second copy of Γ_0 corresponding to t. Then $t_{2,1} \in \Gamma^*_1$ is the vertex corresponding to t_2, which belongs the first copy of the $\gamma_n(\Gamma_0; s; D)$. Put $R'_1 = f'_2$, where f'_2 is the edge of the graph Γ^*_1 labelled $\mathrm{Lab}(f_2)$ and starting at the vertex $t_{2,1}$.

Consider the case $k > 0$. Suppose that f is the edge of S next to $\omega(R_k)$, where S is a maximal u_1-cycle containing R_k. If $\mathrm{Lab}(f) \in A$ then D denotes A, if $\mathrm{Lab}(f) \in B$ then D denotes B. Put $q = \omega(f)$. Consider the graph $\Gamma^*_{k+1} = \gamma_n(\Gamma_k; q; D)$. Put $R'_{k+1} = R_{k,1} \cup f_1$, where the path $R_{k,1}$ and the edge f_1 belong to the first copy of Γ_k in Γ^*_{k+1} and correspond to R_k and f.

Consider a graph X, satisfying Properties (1),(2). Suppose that p is a vertex of X and each u_1-cycle in X has no close edges, then, in the graph $\gamma_r(X; p; K)$, where K is A or B, each u_1-cycle V is the union of r copies of a u_1-cycle W of the graph X. This means that, if W has no close edges, then V has no close edges too. Let $\mathrm{l}(V)$ and $\mathrm{l}(W)$ be the lengths of the u_1-cycles V and W respectively. Since V and W have no close edges $\mathrm{l}(V)=r\mathrm{l}(W)$. Owing to these properties, we conclude that the graph Γ^*_{k+1} satisfies Property (5), and since $\Gamma^*_{k+1} = \gamma_n(Y; z; K)$, n and length of each u_i-cycle of the graph Y for $i = 1, \dots, m$ are powers of p Property (4) is also fulfilled for the graph Γ^*_{k+1}.

The length of S becomes n times greater, and there exists a u_j-cycles in Γ_k, whose length also becomes n times greater in Γ^*_{k+1} for each $j = 2, ..., m$. Properties (4) and (5) shows that, if there exists a u_j-cycle in Γ^*_{k+1}, whose length coincide with the length of a u_1-cycle which is maximal in Γ^*_{k+1}, then this u_j-cycle must contain the path R'_{k+1}. So if we suppose that Condition (3) is fulfilled for Γ^*_{k+1}, then R'_{k+1} is contained in all maximal u_j-cycles of Γ^*_{k+1} and, if this is true for each $j = 2, ..., m$, then this graph satisfies Property (6), and we put $\Gamma_{k+1} = \Gamma^*_{k+1}$ and $R_{k+1} = R'_{k+1}$. Since u_1, u_j do not belong to conjugate cyclic subgroups for each $j = 2, ..., m$, there exists a number k such that the graph Γ^*_{k+1} does not satisfy Property (3). In this case Property (4) shows that the least common multiple of the lengths of all u_1-cycles and the least common multiple of the lengths of all u_j-cycles in Γ^*_{k+1} are different for some $j = 2, ..., m$. So if φ_1 is the homomorphism of the group G which corresponds to the action of G on the set of vertices of the graph Γ^*_{k+1} then $\varphi_1(G)$ is a finite group and

$\varphi_1(u_1), ..., \varphi_1(u_m)$ are nonidentity p-elements and $\{u_1, ..., u_m\} = \alpha \cup \beta$, where α and β are nonempty sets with empty intersection, moreover, for each $u_i \in \alpha$ and $u_j \in \beta$, we have $|\varphi_1(u_i)| > |\varphi_1(u_j)|$.

If $m = 2$ the proof of this statement shows that lemma 3 is true for the case $m = 2$. So we may apply the induction on m.

The set $\{u_1, ..., u_m\}$ can be presented as the union of two disjoint subsets $\{z_1, ..., z_s\}$ and $\{y_1, ..., y_r\}$ and, for each i and j such that $1 \leqslant i \leqslant s$ and $1 \leqslant j \leqslant r$, we have $|\varphi_1(z_i)| > |\varphi_1(y_j)|$. Besides, $\varphi_1(u_1), ..., \varphi_1(u_m)$ are nonidentity p-elements.

Put $\pi' = \pi \cup p$. By the induction hypothesis, there exists a homomorphism φ_2 of group G onto a finite group such that the images of $y_1, ..., y_r$ have pairwise different orders and $p \nmid |\varphi_2(y_i)|$ for each i and each $p \in \pi'$

Suppose that ρ is the set of all prime divisors of the numbers $|\varphi_2(u_1)|, \ldots, |\varphi_2(u_m)|$, $\pi'' = \pi' \cup \rho$. By the induction hypothesis, there exists a homomorphism φ_3 satisfying the requirements for elements $\{z_1, ..., z_s\}$ and the set of primes π''. The homomorphism $\varphi : G \to \varphi_1(G) \times \varphi_2(G) \times \varphi_3(G)$ defined by the formula $\varphi : g \mapsto (\varphi_1(g), \varphi_2(g), \varphi_3(g))$ is as required.

Lemma 4. *Suppose that $G = A * B$, where the groups A and B are finite, w is a nonidentity cyclically reduced element of the Cartesian subgroup C, and the integers $k_1, ..., k_l$ have pairwise different absolute values. Then there exists a homomorphism φ of G onto a finite group such that the orders of the images of the elements $w^{k_1}, ..., w^{k_l}$ are pairwise different.*

Proof.

For each prime divisor p_i of $k_1...k_l$, take a positive integer n_i such that $p_i^{n_i}$ does not divide $k_1...k_l$. According to Lemma 1, there exists a homomorphism φ_{p_i} of G onto a finite group such that $\varphi_{p_i}(w)$ is a p_i-element and $|\varphi_{p_i}(w)| > p_i^{n_i}$. Let $p_1, ..., p_s$ be the set of prime divisors of the numbers $k_1, ..., k_l$. Then the homomorphism $\varphi : G \to \varphi_{p_1}(G) \times ... \times \varphi_{p_s}(G)$ defined by the formula $\varphi : g \mapsto \prod_{i=1}^{s} \varphi_{p_i}(g)$ is as required.

4 Proof of Theorems.

For each homomorphisms φ and ψ of groups A and B respectively we define the homomorphism $\varphi * \psi$ of $A * B$ by the following way. If $s = s_1...s_k \in A * B$ is the normal form of s, then $\varphi * \psi(s) = \phi_1(s_1)...\phi_k(s_k)$, where $\phi_i = \varphi$ in case $s_i \in A$ and $\phi_i = \psi$ if $s_i \in B$.

Proof of Theorems 1 and 2.

Suppose that $G = A * B$ where A and B are either infinite cyclic groups or periodic n-element order separable for each n groups such that $|a| \neq |b|$ for each $a \in A \setminus \{1\}$ and $b \in B \setminus \{1\}$.

Consider the elements $u_1, ..., u_n$ of $A * B$ such that u_i and $u_j^{\pm 1}$ are not conjugate for $i \neq j$

Without loss of generality, we assume that $u_1, ..., u_n$ are cyclically reduced. Consider the following decomposition of the set $\{u_1, ..., u_n\} = \alpha \cup \beta \cup \gamma$, where the sets α, β, and γ are disjoint. The set α consists of all elements of the group A, β consists of all elements of the group B, γ consists of all elements not belonging to $A \cup B$.

We shall prove that there exist homomorphisms φ and ψ of A and B respectively onto finite groups such that the orders of the images of elements of the set $\alpha \cup \beta$ are pairwise different, each element of the set $\alpha \cup \beta \setminus \{1\}$ does not belong to the kernel of φ or ψ, and the images of elements of γ do not belong to groups conjugate to $\varphi(A)$ or $\psi(B)$ after the homomorphism $\varphi * \psi$ of the group G, and the elements $\{\varphi * \psi(u_1), ..., \varphi * \psi(u_n)\}$ are such that $\varphi * \psi(u_i)$ and $\varphi * \psi(u_j^{\pm 1})$ are not conjugate for $i \neq j$.

Consider the set

$$\Lambda = \alpha \cup \Omega \cup (\Omega \cdot \Omega^{-1}),$$

where Ω is the set of all elements of the group A occurring in the normal forms of the elements of γ (if $\Omega = \{\omega_j | j \in J\}$ then $\Omega \cdot \Omega^{-1} = \{\omega_j \omega_k^{-1} | j, k \in J\}$). Next, let us choose a maximal subset Λ' of Λ such that for each different x and y from Λ' x and $y^{\pm 1}$ are not conjugate. Note that $1 \in \Lambda'$.

Consider a homomorphism φ of the group A onto a finite group such that the orders of the images of elements from Λ' are pairwise different.

Similarly, we consider a homomorphism ψ of B for the corresponding elements of this group. Thus, we may assume that A and B are finite (we may consider that for each $a \in \alpha, b \in \beta$, the elements $\varphi * \psi(a)$ and $\varphi * \psi(b)$ have different orders, because of the properties of the free factors). If the set γ is empty use the residual finiteness of the group $A * B$. Suppose that γ is not empty. We may also assume that the set γ lies in the Cartesian subgroup. Indeed, $|G : C| < \infty$. Hence, there exists a positive integer l such that $u^l \in C$ for each $u \in \gamma$. If there exists a homomorphism such that the orders of the images of u^l are pairwise different for each u from γ, then the same is true for the images of the elements u. Furthermore, if $x, y \notin A \cup B$, x and y are cyclically reduced and nonconjugate, then x^n and y^n are nonconjugate for each n (Combinatorial group theory, *Springer-Verlag*, theorem 1.4). Let us decompose the image of the set γ into a union of disjoint subsets $\alpha_1, ..., \alpha_s$, where α_i is a maximal subset in γ such that each element of α_i belongs the cyclic subgroup conjugate to the subgroup generated by some element w_i, and w_i is the same for each element of α_i. We may assume that the set α_i consists of elements $w_i^{k_{ij}}$, and w_i is not a proper power. Let d_i be the greatest common divisor of numbers k_{ij} where i is fixed. Then $w_i^{d_i} \in C$. There exists the number m_i such that $m_i | d_i, w_i^{m_i} \in C$ and if $m_i > 1$ then for each divisor m_i' of m_i which is less than m_i $w_i^{m_i'} \notin C$. Thus we may consider that the set α_i consists of elements $w_i^{k_{ij}}$, $w_i \in C$ and if $w_i = w_i'^{m_i}$, where w_i' is not a proper power then $w_i'^{m_i'} \notin C$ for each m_i' such that $m_i' < m_i, m_i' | m_i$. By Lemma 4, for each α_i, there exists a homomorphism φ_i such that the orders of elements from $\varphi_i(\alpha_i)$ are pairwise different.

The orders of the images of the elements from $\alpha \cup \beta$ after φ_i are pairwise different because $\varphi_i |_A, \varphi_i |_B$ are injective.

The set α_i consists of elements $w_i^{k_{ij}}$. Let π be the set consisting of prime divisors of the orders of elements $\varphi_i(\alpha_i)$ and of the images of α and β under the homomorphism $G \to \varphi_1(G) \times ... \times \varphi_s(G)$ defined by $g \mapsto (\varphi_1(g), ..., \varphi_s(g))$. By Lemma 3 there exists a homomorphism ψ such that the orders of elements $\psi(w_1), ..., \psi(w_s)$ are pairwise different and $p \nmid |\psi(w_i)|$ for each i and for each $p \in \pi$ and $\psi\mid_A, \psi\mid_B$ are injective.

The homomorphism $\chi : G \to \varphi_1(G) \times ... \times \varphi_s(G) \times \psi(G)$ defined by $\chi : g \mapsto (\varphi_1(g), ..., \varphi_s(g), \psi(g))$ is as required.

Proof of Theorem 3.

It is sufficient to show, how to reduce the general case to the case, when the free factors are finite. The proof of this particular case is similar to the proof of Theorems 1 and 2, because after we reduced the general case to this particular case we did not use any properties of finite free factors.

Suppose the elements u, v, w are cyclically reduced and, if x and y are two different elements from $\{u, v, w\}$, then x and $y^{\pm 1}$ are not conjugate.

If none of these elements lies in A, or none of them lies in B, then the proof is the same as the proof of Theorems 1 and 2. Thus, we have to consider two cases: $u \in A \setminus e, v \in B, w \notin A \cup B$ or $u, v \in A, w \in B \setminus e$. In the first case, there exists a homomorphism φ_1 of A onto a finite group such that $\varphi_1(u) \neq e$ and the image of w does not belong to a group conjugate to the image of some free factor after the corresponding homomorphism of the group G, besides, if the order of u is infinite and the order of v is finite, we may assume that $\varphi_1(u^p) \neq e$ for each divisor p of $|v|$. There exists a homomorphism φ_2 of B onto a finite group such that $\varphi_2(v^{|\varphi_1(u)|}) \neq e$ if $v \neq e$. In addition, the image of $\varphi_1 * \mathrm{id}(w)$ does not belong to a group conjugate to the image of a free factor after the corresponding homomorphism of $\varphi_1(A) * B$. If $v = e$, then it is sufficient to satisfy the first condition. In this case, we use the residual finiteness of A and B.

In the second case, if $u \neq e, v \neq e$ there exists a homomorphism φ_1 of A onto a finite group such that $|\varphi_1(u)| \neq |\varphi_1(v)|$ and $\varphi_1(u) \neq e, \varphi_1(v) \neq e$. The homomorphism φ_1 with the above properties exists because the group A is 3-element order separable, and the natural homomorphism $\varphi_2 : \varphi_1(A) * B \to \varphi_1(A)$ is as required. Suppose that $u = e$. If the orders of v and w are finite then $|v|$ and $|w|$ are coprime and we use the residual finiteness of G. Consider the case $u = e$ and the order of v is infinite. Since the group A is residually finite there exists a homomorphism φ_1 of A onto a finite group such that $\varphi_1(v) \neq e$. And if the order of w is finite we may consider that $\varphi_1(v^p) \neq e$ for each divisor p of $|w|$. The group B is residually finite so there exists a homomorphism φ_2 of B onto a finite group such that $\varphi_2(w^{|\varphi_1(v)|}) \neq e$, and we use the residual finiteness of the group $\varphi_1(A) * \varphi_2(B)$.

Acknowledgements

The author thanks A. A. Klyachko for valuable comments and setting the problem.

References

1. Endimioni, G. Elements with the same normal closure in a metabelian group. *The Quarterly Journal of Mathematics* **58(1)**: 23-29, 2006.

2. Kargapolov, M. I., Merzlyakov, Yu. I. (1977). Foundations of group theory. *Nauka*.

3. Klyachko, A. A. Equations over groups, quasivarieties, and a residual property of a free group. *Journal of group theory* **2**: 319-327, 1999.

4. Lyndon, R. C., Schupp, P. E. (1977). Combinatorial group theory. *Springer-Verlag*.

5. Yedynak, V. V. Separability with respect to order. *Vestnik Mosk. Univ. Ser. I Mat. Mekh.* **3**: 56-58, 2006.

Yedynak V. V., a graduate student of Moscow State Univ.

Yedynak V. V., Trudovaya street 14-b, 32, Ivanteevka, Moscow Region, 141281, Russia.